



Research note

First record of *Spathoteredo spatha* (Mollusca: Teredinidae) in Venezuela

Primer registro de Spathoteredo spatha (Mollusca: Teredinidae) en Venezuela

Marcel Velásquez* and Irelis López

Facultad Ciencias del Mar, Universidad Católica del Norte, Larrondo 1281, Coquimbo, Chile

Received 7 April 2014; accepted 2 September 2014

Abstract

We document the first record of shipworm mollusk *Spathoteredo spatha* (Jeffreys, 1860) in Venezuelan coastal waters. Also, we present data about the distribution of the species in the Caribbean Sea.

All Rights Reserved © 2015 Universidad Nacional Autónoma de México, Instituto de Biología. This is an open access item distributed under the Creative Commons CC License BY-NC-ND 4.0.

Keywords: Teredinids; Shipworms; Caribbean Sea

Resumen

Documentamos el primer registro del molusco perforador de madera *Spathoteredo spatha* (Jeffreys, 1860) en aguas de las costas de Venezuela. Además, se presentan algunos datos de la distribución de esta especie en el mar Caribe.

Derechos Reservados © 2015 Universidad Nacional Autónoma de México, Instituto de Biología. Este es un artículo de acceso abierto distribuido bajo los términos de la Licencia Creative Commons CC BY-NC-ND 4.0.

Palabras clave: Teredinidos; Moluscos perforadores; Mar Caribe

There are currently 72 known species of the Teredinidae family, and these are distributed throughout the seas and oceans around the world (Bouchet, 2013; Turner, 1966). Fourteen species of this family have been reported in the Caribbean Sea (Abbott, 1974; Díaz & Puyana, 1994; Miloslavich et al., 2010; Turner, 1966), of which 11 have been recorded in the Venezuelan coast (Ewald, Severejn, & Espinoza, 1984; Nair, 1975; Turner, 1954; 1955; 1966). There have been no recorded observations of new teredinids in Venezuela since the last study of boring molluscs done by Ewald et al. (1984) in the Maracaibo Lake. Here *Spathoteredo spatha* is recorded for the first time in Venezuela. This observation also represents the first record of a Teredinidae species in this country in 30 years.

Material examined

A total of 17 live specimens of *S. spatha* were collected between October 2007 and February 2014, in Punta de Piedras

Lagoon, Margarita Island, Venezuela. All of them were found boring into sunken wood of *Rhizophora mangle*. The samples were preserved in ethanol 70%, with some being deposited to the Malacology collection at Oceanology Museum Hno. Benigno Román of the Marine Station Research of Margarita (EDIMAR), Margarita Island, Venezuela, under the Catalog number MOBR-M-3985. Also, for the correct identification of our samples these were compared with specimens of the Malacology Collection at Museum of Comparative Zoology (MCZ), Harvard University, Cambridge, Massachusetts.

Spathoteredo spatha (Jeffreys, 1860)

Description

Shell with a heavy brown periostracum; valves small, globose, anterior face with numerous denticulate ridges, typical of Teredinidae (Turner, 1966). Pallets are oval with a band of periostracum on the median area of the outer face and cover a small part of the internal face. The upper portions of the blade covered with a calcareous deposit, segmented having a ribbed

* Corresponding author.

E-mail address: marcelvelasquez2@gmail.com (M. Velásquez).




Figure 1. Pallets of *Spathoteredo spatha*. A, internal face; B, outer face. Scale= 1 mm.

appearance (Fig. 1). The size of the pallets varied between 2.38-3.96 mm. The siphons are united except at the tip. All specimens were collected from sunken wood of *Rhizophora mangle* between 1.5-2 meters deep.

Comparative material

Indian River, Florida, USA (MCZ-356880); Puerto Cortés, Honduras (MCZ-227262); Fajardo, Puerto Rico (MCZ-170703); Galeta Island, Panama (MCZ-356879); Santa Marta, Chengue, Colombia (MCZ-278056).

Geographical distribution

Atlantic North (West coast), USA (Turgeon et al., 1998); Atlantic North (East coast): United Kingdom, England (Turner, 1966); Gulf of Mexico and Caribbean Sea (Turner, 1966).

The specimens examined are morphologically similar to the descriptions of Turner (1966) and the specimens which were compared from the Museum of Comparative Zoology. However, these show some differences with the illustration pallets of *S. spatha* presented by Turner (1966), since the figure shows a whitish papillose calcareous deposit in the upper portion of the blade and this feature is not present in the pallets of our specimens, nor in the majority of comparative specimens from the museum.

In Teredinidae species it is common to find high intraspecific variation that is evident in the morphology of the pallets and valves, which are the principal structures used for the identification of the species (Turner, 1966). This variation is a product of the boring life style and environmental conditions (Borges, Merckelbach, Sampaio, & Cragg, 2014; Cragg, Jumel, Al-Horani, & Hendi, 2009) and this was generated as a result of the erroneous identifications in the past (Borges et al., 2014); a fact that can be demonstrated by the number of existent synonymy (Turner, 1966).

Spathoteredo spatha was described for the first time by Jeffreys (1860), and named *Teredo spatha*, this description was based on specimens obtained from driftwood from Guernsey's coast, England. Later, Roch (1931) proposed a new name (*Teredo molli*) for specimens of *S. spatha* collected from driftwood in the Atlantic Ocean. The organization of the species into the currently known genus *Spathoteredo*, which consists of the 2 species *S. spatha* and *S. obtusa*, did not occur until 1966 when R. Turner proposed this in the book *The Survey Illustrated Catalogue of the Teredinidae*. The first species has a distribution range principally from Caribbean Sea to both sides of the North Atlantic and the South Atlantic waters. However, there is a report of *S. spatha* in European waters (Gofas, 2013; Gofas, Le Renard, & Bouchet, 2001), but this is not considered an established species for this zone because all specimens were found either in driftwood carried mainly by the Gulf Stream to the British Isles or in driftwood in the Mediterranean (Borges et al., 2014). The second species is distributed throughout the Pacific Ocean (Africa (East coast), Madagascar, Red Sea, Persian Gulf, Pacific Islands, Southeast Asia, Indonesia, New Guinea, Philippine Islands) (Turner, 1966).

The collection of *S. spatha* specimens was carried out in 2 different years in the locality of Punta de Piedras Lagoon, so it can therefore be concluded that this species is established in this locality.

The principal author is grateful to Adam Baldinger, Curatorial Associate of Malacological Collection of the Comparative Zoology Museum of Harvard University (MCZ), for allowing the review of the Teredinidae samples of the collection in order to make comparisons with the specimens used in this study. Thanks also to Alex Carter for his helpful comments on the translation of the manuscript.

References

- Abbott, R. T. (1974). *American Seashells* (2nd ed.). New York: Van Nostrand Reinhold Company.
- Borges, L., Merckelbach, L., Sampaio, I., & Cragg, S. (2014). Diversity, environmental requirements, and biogeography of bivalve wood-borers (Teredinidae) in European coastal waters. *Frontiers in Zoology*, 11, 13.
- Bouchet, P. (2013). Teredinidae. Accessed through: *World register of marine species*. Retrieved on June 4th, 2014 from <http://www.marinespecies.org/aphia.php?p=taxdetails&id=253>
- Cragg, S., Jumel, M., Al-Horani, F., & Hendi, I. (2009). The life history characteristics of the wood-boring *Teredo bartschi* are suited to the elevated salinity, oligotrophic circulation in the Gulf of Aqaba, Red Sea. *Journal of Experimental Marine Biology and Ecology*, 375, 99–105.
- Díaz, J. M., & M. Puyana. (1994). *Moluscos del Caribe Colombiano*. Un catálogo ilustrado. Bogotá: Colciencias, Fundación Natura e INVEMAR.
- Ewald, J. J., Severeyn, H. & Espinoza, D. (1984). La fauna acuática de invertebrados de la Cuenca del Lago de Maracaibo: I. Los moluscos bivalvos perforadores. *Ciencias, Revista Científica de la Universidad del Zulia*, 1(2), 41–64.
- Gofas, S. (2013). *Spathoteredo spatha* (Jeffreys, 1860). Accessed through: *World register of marine species*. Retrieved on March 31st, 2014 from <http://www.marinespecies.org/aphia.php?p=taxdetails&id=141604>
- Gofas, S., Le Renard, J., & Bouchet, P. (2001). Mollusca. En M. J. Costello, C. Embrow & R. White (Eds.), *European register of marine species: a check-list of the marine species in Europe and a bibliography of guides to their identification* (pp. 180–213). Collection Patrimoine Naturels 50.

- Jeffreys, J. G. (1860). A synoptical list of the British species of *Teredo* with a notice of the exotic species. *Annals and Magazine of Natural History*, 6, 121–127.
- Miloslavich, P., Díaz, J., Klein, E., Alvarado, J., Díaz, C., Gobin, J. et al. (2010). Marine biodiversity in the Caribbean: regional estimates and distribution patterns. *PLoS ONE*, 5, 1–25.
- Nair, N. B. (1975). Shipworms of Venezuela report on a collection from the gulf of Cariaco. *Boletín del Instituto Oceanográfico de Venezuela*, 14, 129–146.
- Roch. F. (1931). Die Terediniden der Skandinavischen. *Museum Sammlung Arkiv för Zoology*, 22 (13), 1–29.
- Turgeon, D. D., Quinn Jr, J. F., Bogan, A. E., Coan, E. V., Hochberg, F. G., Lyons, W. C. et al. (1998). *Common and scientific names of aquatic invertebrates from the United States and Canada: mollusks* (2nd ed.). Bethesda: American Fisheries Society (Special publication 26).
- Turner, R. D. (1954). The family Pholadidae in the western Atlantic and the eastern Pacific, Part I: Pholadinae. *Johnsonia*, 3, 1–63.
- Turner, R. D. (1955). The family Pholadidae in the western Atlantic and the eastern Pacific, Part II: Martesiinae, Juannetiinae and Xylophaginae. *Johnsonia*, 3, 65–100.
- Turner, R. (1966). *A survey and illustrated catalogue of the Teredinidae (Mollusca: Bivalvia)*. Cambridge: The Museum of Comparative Zoology, Harvard University.